

IMO Shortlist 2008

Berlin, Germany

Algebra

- 1 Find all functions $f : (0, \infty) \mapsto (0, \infty)$ (so f is a function from the positive real numbers) such that

$$\frac{(f(w))^2 + (f(x))^2}{f(y^2) + f(z^2)} = \frac{w^2 + x^2}{y^2 + z^2}$$

for all positive real numbers w, x, y, z , satisfying $wx = yz$.

Author: Hojoo Lee, South Korea

- 2 (i) If x, y and z are three real numbers, all different from 1, such that $xyz = 1$, then prove that $\frac{x^2}{(x-1)^2} + \frac{y^2}{(y-1)^2} + \frac{z^2}{(z-1)^2} \geq 1$. (With the \sum sign for cyclic summation, this inequality could be rewritten as $\sum \frac{x^2}{(x-1)^2} \geq 1$.)

(ii) Prove that equality is achieved for infinitely many triples of rational numbers x, y and z .

Author: Walther Janous, Austria

- 3 Let $S \subseteq \mathbb{R}$ be a set of real numbers. We say that a pair (f, g) of functions from S into S is a *Spanish Couple* on S , if they satisfy the following conditions: (i) Both functions are strictly increasing, i.e. $f(x) < f(y)$ and $g(x) < g(y)$ for all $x, y \in S$ with $x < y$;

(ii) The inequality $f(g(g(x))) < g(f(x))$ holds for all $x \in S$.

Decide whether there exists a Spanish Couple on the set $S = \mathbb{N}$ of positive integers; [/*:m] on the set $S = \{a - \frac{1}{b} : a, b \in \mathbb{N}\}$ [/*:m]

Proposed by Hans Zantema, Netherlands

- 4 For an integer m , denote by $t(m)$ the unique number in $\{1, 2, 3\}$ such that $m + t(m)$ is a multiple of 3. A function $f : \mathbb{Z} \rightarrow \mathbb{Z}$ satisfies $f(-1) = 0$, $f(0) = 1$, $f(1) = -1$ and $f(2^n + m) = f(2^n - t(m)) - f(m)$ for all integers $m, n \geq 0$ with $2^n > m$. Prove that $f(3p) \geq 0$ holds for all integers $p \geq 0$.

Proposed by Gerhard Woeginger, Austria

- 5 Let a, b, c, d be positive real numbers such that $abcd = 1$ and $a + b + c + d > \frac{a}{b} + \frac{b}{c} + \frac{c}{d} + \frac{d}{a}$. Prove that

$$a + b + c + d < \frac{b}{a} + \frac{c}{b} + \frac{d}{c} + \frac{a}{d}$$

Proposed by Pavel Novotn, Slovakia

IMO Shortlist 2008

Berlin, Germany

- 6] Let $f : \mathbb{R} \rightarrow \mathbb{N}$ be a function which satisfies $f\left(x + \frac{1}{f(y)}\right) = f\left(y + \frac{1}{f(x)}\right)$ for all $x, y \in \mathbb{R}$.
Prove that there is a positive integer which is not a value of f .

Proposed by ymantas Darbnas (Zymantas Darbenas), Lithuania

- 7] Prove that for any four positive real numbers a, b, c, d the inequality

$$\frac{(a-b)(a-c)}{a+b+c} + \frac{(b-c)(b-d)}{b+c+d} + \frac{(c-d)(c-a)}{c+d+a} + \frac{(d-a)(d-b)}{d+a+b} \geq 0$$

holds. Determine all cases of equality.

Author: Darij Grinberg (Problem Proposal), Christian Reiher (Solution), Germany

IMO Shortlist 2008

Berlin, Germany

Combinatorics

- 1 In the plane we consider rectangles whose sides are parallel to the coordinate axes and have positive length. Such a rectangle will be called a *box*. Two boxes *intersect* if they have a common point in their interior or on their boundary. Find the largest n for which there exist n boxes B_1, \dots, B_n such that B_i and B_j intersect if and only if $i \not\equiv j \pm 1 \pmod{n}$.

Proposed by Gerhard Woeginger, Netherlands

- 2 Let $n \in \mathbb{N}$ and A_n set of all permutations (a_1, \dots, a_n) of the set $\{1, 2, \dots, n\}$ for which

$$k|2(a_1 + \dots + a_k), \text{ for all } 1 \leq k \leq n.$$

Find the number of elements of the set A_n .

- 3 In the coordinate plane consider the set S of all points with integer coordinates. For a positive integer k , two distinct points $a, B \in S$ will be called *k-friends* if there is a point $C \in S$ such that the area of the triangle ABC is equal to k . A set $T \subset S$ will be called *k-clique* if every two points in T are *k-friends*. Find the least positive integer k for which there exists a *k-clique* with more than 200 elements.

Proposed by Jorge Típe, Peru

- 4 Let n and k be positive integers with $k \geq n$ and $k - n$ an even number. Let $2n$ lamps labelled $1, 2, \dots, 2n$ be given, each of which can be either *on* or *off*. Initially all the lamps are off. We consider sequences of steps: at each step one of the lamps is switched (from on to off or from off to on).

Let N be the number of such sequences consisting of k steps and resulting in the state where lamps 1 through n are all on, and lamps $n + 1$ through $2n$ are all off.

Let M be number of such sequences consisting of k steps, resulting in the state where lamps 1 through n are all on, and lamps $n + 1$ through $2n$ are all off, but where none of the lamps $n + 1$ through $2n$ is ever switched on.

Determine $\frac{N}{M}$.

Author: Bruno Le Floch and Ilia Smilga, France

- 5 Let $S = \{x_1, x_2, \dots, x_{k+l}\}$ be a $(k + l)$ -element set of real numbers contained in the interval $[0, 1]$; k and l are positive integers. A k -element subset $A \subset S$ is called *nice* if

$$\left| \frac{1}{k} \sum_{x_i \in A} x_i - \frac{1}{l} \sum_{x_j \in S \setminus A} x_j \right| \leq \frac{k+l}{2kl}$$

IMO Shortlist 2008

Berlin, Germany

Prove that the number of nice subsets is at least $\frac{2}{k+l} \binom{k+l}{k}$.

Proposed by Andrey Badzhan, Russia

- 6 For $n \geq 2$, let S_1, S_2, \dots, S_{2^n} be 2^n subsets of $A = \{1, 2, 3, \dots, 2^{n+1}\}$ that satisfy the following property: There do not exist indices a and b with $a < b$ and elements $x, y, z \in A$ with $x < y < z$ and $y, z \in S_a$, and $x, z \in S_b$. Prove that at least one of the sets S_1, S_2, \dots, S_{2^n} contains no more than $4n$ elements.

Proposed by Gerhard Woeginger, Netherlands

IMO Shortlist 2008

Berlin, Germany

Geometry

- 1] Let H be the orthocenter of an acute-angled triangle ABC . The circle Γ_A centered at the midpoint of BC and passing through H intersects the sideline BC at points A_1 and A_2 . Similarly, define the points B_1, B_2, C_1 and C_2 .

Prove that six points A_1, A_2, B_1, B_2, C_1 and C_2 are concyclic.

Author: Andrey Gavriluk, Russia

- 2] Given trapezoid $ABCD$ with parallel sides AB and CD , assume that there exist points E on line BC outside segment BC , and F inside segment AD such that $\angle DAE = \angle CBF$. Denote by I the point of intersection of CD and EF , and by J the point of intersection of AB and EF . Let K be the midpoint of segment EF , assume it does not lie on line AB . Prove that I belongs to the circumcircle of ABK if and only if K belongs to the circumcircle of CDJ .

Proposed by Charles Leytem, Luxembourg


- 3] Let $ABCD$ be a convex quadrilateral and let P and Q be points in $ABCD$ such that $PQDA$ and $QPBC$ are cyclic quadrilaterals. Suppose that there exists a point E on the line segment PQ such that $\angle PAE = \angle QDE$ and $\angle PBE = \angle QCE$. Show that the quadrilateral $ABCD$ is cyclic.

Proposed by John Cuya, Peru

- 4] In an acute triangle ABC segments BE and CF are altitudes. Two circles passing through the point A and F and tangent to the line BC at the points P and Q so that B lies between C and Q . Prove that lines PE and QF intersect on the circumcircle of triangle AEF .

IMO Shortlist 2008

Berlin, Germany



[Diagram by azjps]

Proposed by Davood Vakili, Iran

- 5] Let k and n be integers with $0 \leq k \leq n - 2$. Consider a set L of n lines in the plane such that no two of them are parallel and no three have a common point. Denote by I the set of intersections of lines in L . Let O be a point in the plane not lying on any line of L . A point $X \in I$ is colored red if the open line segment OX intersects at most k lines in L . Prove that I contains at least $\frac{1}{2}(k+1)(k+2)$ red points.

Proposed by Gerhard Woeginger, Netherlands

- 6] There is given a convex quadrilateral $ABCD$. Prove that there exists a point P inside the quadrilateral such that

$$\angle PAB + \angle PDC = \angle PBC + \angle PAD = \angle PCD + \angle PBA = \angle PDA + \angle PCB = 90^\circ$$

if and only if the diagonals AC and BD are perpendicular.

Proposed by Dukan Dukic, Serbia

- 7] Let $ABCD$ be a convex quadrilateral with BA different from BC . Denote the incircles of triangles ABC and ADC by k_1 and k_2 respectively. Suppose that there exists a circle k tangent to ray BA beyond A and to the ray BC beyond C , which is also tangent to the lines AD and CD .

Prove that the common external tangents to k_1 and k_2 intersect on k .

Author: Vladimir Shmarov, Russia

IMO Shortlist 2008

Berlin, Germany

Number Theory

- 1] Let n be a positive integer and let p be a prime number. Prove that if a, b, c are integers (not necessarily positive) satisfying the equations

$$a^n + pb = b^n + pc = c^n + pa$$

then $a = b = c$.

Proposed by Angelo Di Pasquale, Australia

- 2] Let a_1, a_2, \dots, a_n be distinct positive integers, $n \geq 3$. Prove that there exist distinct indices i and j such that $a_i + a_j$ does not divide any of the numbers $3a_1, 3a_2, \dots, 3a_n$.

Proposed by Mohsen Jamaali, Iran

- 3] Let a_0, a_1, a_2, \dots be a sequence of positive integers such that the greatest common divisor of any two consecutive terms is greater than the preceding term; in symbols, $\gcd(a_i, a_{i+1}) > a_{i-1}$. Prove that $a_n \geq 2^n$ for all $n \geq 0$.

Proposed by Morteza Saghafian, Iran

- 4] Let n be a positive integer. Show that the numbers

$$\binom{2^n - 1}{0}, \binom{2^n - 1}{1}, \binom{2^n - 1}{2}, \dots, \binom{2^n - 1}{2^{n-1} - 1}$$

are congruent modulo 2^n to $1, 3, 5, \dots, 2^n - 1$ in some order.

Proposed by Duskan Dukic, Serbia

- 5] For every $n \in \mathbb{N}$ let $d(n)$ denote the number of (positive) divisors of n . Find all functions $f : \mathbb{N} \rightarrow \mathbb{N}$ with the following properties: $d(f(x)) = x$ for all $x \in \mathbb{N}$. [*/:m] $f(xy)$ divides $(x-1)y^{xy-1}f(x)$ for all $x, y \in \mathbb{N}$. [*/:m]

Proposed by Bruno Le Floch, France

- 6] Prove that there are infinitely many positive integers n such that $n^2 + 1$ has a prime divisor greater than $2n + \sqrt{2n}$.

Author: Kestutis Cesnavicius, Lithuania